

# Driven motion of the third phase on the oil | water interface by three-phase boundary reactions

Jingyuan Chen, Masanori Satoh, Tomohiro Imamura, Hirokazu Toda, Koichi Aoki

Department of Applied Physics, University of Fukui,  
3-9-1 Bunkyo, Fukui, 910-8507 Japan

A highly competitive system is the reaction at the three-phase boundary at which the electrode was exposed to both the oil and the water phase, explored by Scholz and coworkers<sup>1-3</sup>. It is composed of an oil droplet mounted on the electrode in an aqueous solution. A typical cell apparatus is composed of the oil droplet including only a redox species, which is located on an electrode in an aqueous solution including supporting electrolyte<sup>4-7</sup>, as is shown in Fig.1A. An air bubble mounted on an oil | water interface is a probe not only of interfacial tension but also of properties of three-phase boundary reactions<sup>8</sup>.

This report is devoted to investigating the static and dynamic behavior of the air bubble on the O|W interface in the context of the three-phase boundary reaction of ferrocene. The bubble kept spontaneously moving on the oil|water interface without applying any force, when it is less than 0.3 mm in diameter. The motion was irregular without decay, suggesting a contribution of Brownian motion. When the three-phase boundary reaction occurred, the bubble motion stopped. It was observed only in the potential domain at which ferrocene is oxidized. The force balance of the air bubble at the O|W interface will be formulated in terms of the interfacial tensions and the buoyancy. It will be demonstrated that the electrode reaction varies the surface tension at the O|W interface to cause surface convection.




Fig.1. Illustration (A) of a hemispherical oil droplet including an air bubble mounted on an electrode in aqueous solution, and interpretation (B) for the three-phase boundary reaction of ferrocene (Fc) in the oil phase. Supporting ions, M<sup>+</sup> and A<sup>-</sup>, form a double layer.

## References

1. F. Scholz, S. Komorsky-Lovric, M. Lovric, *Electrochem. Commun.*, **2000**, 2, 112.
2. M. Donten, Z. Stojek, F. Scholz, *Electrochem. Commun.*, **2002**, 4, 324.
3. M. Lovric, F. Scholtz, *J. Solid State Electrochem.*, **1997**, 1, 108.
4. J.Chen, M.Sato, *J. Electroanal. Chem.*, **2004**, 572, 153
5. P.Tasakorn, J.Chen, K.Aoki, *J. Electroanal. Chem.*, **2002**, 533, 119.
6. K.Aoki, P.Tasakorn, J.Chen, *J. Electroanal. Chem.*, **2003**, 542, 51
7. K.Aoki, M.Satoh, J.Chen, T.Nishiumi, *J. Electroanal. Chem.*, **2006**, 598, 103.
8. M.Satoh, K.Aoki, J.Chen, *Langmuir*, **2008**, 24(8), 4364.